

X-TreemTM Moly



Reg. No. B 4865 Act No. 36 of 1947 / Wet Nr. 36 van 1947

A soluble crystalline powder sodium-molybdate for the supply of molybdenum to crops.

FERTILIZER GROUP 2

STORE AT ROOM TEMPERATURE. KEEP OUT OF DIRECT SUNLIGHT.

Active Ingredients	
Molybdenum	394 g/kg

Registration Holder N Laboratories (PTY) LTD
Reg 2015/165672/07 • No. 144, 2nd Avenue,
Modder East Orchards, Delmas, 2210
Tel: +27 (0) 82 738 0080 • Fax: +27 (0) 86 547 6711

Marketed and Distributed by: MBFi (PTY) LTD.
REG No 2013/211882/07
P.O Box 1137, Delmas, Mpumalanga, 2210
orders@mbfi.co.za • www.mbfi.co.za

Date of manufacture:
Datum van vervaardiging:

Batch No:
Lotnommer:

1kg

X-Treem is a Registered Trademark of N Laboratories (PTY) LTD

X-Treem^{Moly}

Reg. No. B 4865 Act No. 36 of 1947 / Wet Nr. 36 van 1947

A soluble crystalline powder sodium-molybdate for the supply of molybdenum to crops.

FERTILIZER GROUP 2

STORE AT ROOM TEMPERATURE. KEEP OUT OF DIRECT SUNLIGHT

ACTIVE INGREDIENTS:

Molybdenum (Mo)* 394 g/kg

Net weight1kgNetto

Registration Holder / Registrasiehouer:

N Laboratories (PTY) LTD
Reg. 2015/165672/07
No. 144, 2nd Avenue, Modder East Orchards, Delmas, 2210
Tel: +27 (0) 82 736 0080 • Fax: +27 (0) 86 547 6711

Marketed and Distributed by:

MBFI (PTY) LTD
Reg No. 2013/211882/07
P.O Box 1137, Delmas, Mpumalanga, 2210
Email: orders@mbfi.co.za • Web: www.mbfi.co.za

X-Treem is a Registered Trademark of N Laboratories (PTY) LTD

DIRECTIONS FOR USE - USE ONLY AS INDICATED

Use exclusively as fertilizer. X-Treem Moly is a crystalline powder of 100% Na-molybdate, which may be used either as a seed dressing or a foliar spray to combat molybdenum deficiency in agricultural crops.

MIXING AND APPLICATION DETAILS:

X-Treem Moly should preferably be applied in the early morning or late afternoon. The rate of application is dependant on the soil fertility and reaction required. The lower rates should be used for maintenance while the higher rates will correct nutritional deficiencies.

STORAGE:

X-Treem Moly should be stored out of direct sunlight and in a dry and cool place under frost free conditions. Keep the products locked up and out of the reach of children and pets. Avoid contamination of foodstuff or water sources.

CROPS	DOSAGE/HA	REMARKS
Lucerne	125 to 250 g/ha	Seed treatment simultaneously with inoculation
Grass and Clover Pastures	155 g/ha	Foliar spray onto young growth pastures after cutting.
Tomatoes	250 g/ha	Molybdenum deficiency occurs generally in the Lowveld region, especially on acid soils. Apply to the planting furrow just before transplanting or as a foliar spray.
Maize	60 g/1.25L of water	Place 100kg of seed in a drum with a watertight lid. Add the solution and turn the drum over for 15 to 20 minutes by rolling or turning on an axle by means of a handle. The seed may be treated any time before planting.
Cruciferous Crops	100 to 250 g/ha	Spray the plants with a solution in water.
Curcubits	2 g/L water	Let the seed soak overnight in a 0.2% solution and plant directly afterwards.
Soybean	50 to 150 g/ha	Seed treatment simultaneously with inoculation
Sunflowers	25 g/25kg seed 5 g/25kg seed	A solution of the trace element should be applied uniformly to the seed. Apply to the seedlings as a full cover foliar spray.

REPEAT TREATMENTS:

Depending on the quantity of X-Treem Moly applied, it should not be necessary to treat again for periods listed below:

Quantity required per ha	Period for which no further treatment will be necessary
More than 150g	4 to 5 years
75 to 150 g	2 to 3 years
< 75 g	1 year