



# NITRO-LIQ

## SOYBEAN



Reg No: L 8986 Act No. 36 of 1947

(Bacterial legume Inoculant for the effective nitrogen fixation on soybean (*Glycine max*))

Active ingredient:

Live legume bacteria for soybeans

(*Bradyrhizobium japonicum*) Minimum  $2 \times 10^9$  cfu/ml

### SHAKE BEFORE USING

Batch No:

Expiry Date:



Leaders in Science. Partners in Growth.

Net Content  
**150ml**  
Treats up to  
50 kg of seed

Developed and Registered by: Natrol (PTY) LTD • Reg 2015/165672/07 • No. 144, 2nd Avenue, Modder East Orchards, Delmas, 2210. • Tel: +27 (0) 82 738 0080 • Fax: +27 (0) 86 547 6711

Produced by: Rizobacter S.A. SENASA N°: 21193 • Laboratory Reg: PB0013

Pergamino (Bs.As.) Argentina • Marketed and Distributed by: MBFi (PTY) LTD  
Reg 2013/21 1882/07 • P.O Box 1137, Delmas, 2210 • orders@mbfi.co.za • www.mbfi.co.za

## Warning

1. Store in a cool (4-24°C), dark, dry place.
2. Do not use after expiry date.
3. Do not store near to pesticides.
4. Do not use for any other legume except as indicated.
5. Use immediately once the sachet has been opened.
6. Not harmful to humans and animals.
7. Once the seed has been treated with the Inoculant, plant as soon as manageable.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions; quality of dilution water; compatibility with other substances not indicated on the label, as well as by the method, time and accuracy of application. Furthermore the registration holder, does not accept responsibility for damage to crops, vegetation, the environment, or harm to man or animal, or for lack of performance of the remedy concerned, due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

### Direction for use:

Use only as indicated on this label. Use double recommended dosage under poor planting conditions or when thiram containing preparations are used.

Application method: Seed treatment or Infurrow	Dosage / 100 kg of seed	Remarks
Seed treatment: Nitro-Liq Soybean	300ml	Apply to the seed and mix thoroughly until it is evenly coated.
Infurrow: Nitro-Liq Soybean	300-450 ml per hectare	Mix with water and apply infurrow directly on the seed

**Note: Please ensure the seed is dry before planting starts. Use talc/graphite to shorten this interval.**

**PreMax Protector:** We recommend the use of PreMax protector with the Nitro-Liq Soybean Inoculant which will help to increase bacterial survival on the seed or in the soil under most stress conditions. Premix the PreMax Protector and the Nitro-Liq Soybean Inoculant and then add to the seed and mix thoroughly until it is evenly coated. Plant treated seed as soon as manageable.

**Note:** The seed quality, size, moisture content and germination percentage prior to treatment will affect the seed treatment load quantity and consistency of the seed after treatment. Please ensure that seed quality has been tested prior to treatment. This product was tested on registered and certified seed; and recommendations are made accordingly to ensure quality in seed treatment performance and planting practices. High humidity may result in seed becoming sticky after treatment. If this occurs, or if the time between mixing and planting needs to be shortened, follow planter recommendations for the use of talc and graphite to dry seed out. If the application of talc/graphite is not mentioned in planter recommendations, add as required or until seed is sufficiently dry.

# N-Factor 1



NITROLS0001

NitroLiq Soybean 150ml

