

SIGNAL[®] PRO

Crop Defence System

Reg. No B 6697 Act No. 36 of 1947 / Reg. Nr. B 6697 Wet Nr. 36 van 1947

READ THE LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN AND ANIMALS

Group/Groep

2

Fertilizer/Kunsmis

A LIQUID FERTILIZER COMPLEX COMBINATION OF MICRONUTRIENTS (COPPER, MANGANESE AND ZINC) FOR CROPS AS LISTED.

Danger



Hazard Statements:

- Causes severe skin burns and eye damage.
- Toxic to aquatic life with long lasting effects.

Precautionary Statements:

- Do not breathe mist and spray.
- Wash hands thoroughly after handling.
- Avoid release to the environment.

Composition:

Copper (Cu)*	3.2 g/kg	4.0 g/L	Potassium (K)	9.7 g/kg	12.2 g/L
Manganese (Mn)*	23.0 g/kg	28.7 g/L	Phosphorus (P)	6.5 g/kg	8.12 g/L
Zinc (Zn)*	35.2 g/kg	44.0 g/L			

*Organically Chelated
Contains Salicylic Acid
S.G (at 20°C): 1.25

UN 1760

Net Volume:
20 L



Registered, Marketed and Distributed by / Geregistreer, Bemark en Versprei deur:
MBFi (PTY) LTD • Reg. No.: 2015/165672/07

No. 144 Avenue 2, Modder East Orchards, Delmas, 2210
Tel: +27 (0) 82 738 0080 • Email: orders@mbfi.co.za • Web: www.mbfi.co.za

IN CASE OF POISONING, CALL THE FOLLOWING EMERGENCY NUMBER / IN GEVAL VAN
VERGIFTIGING, SKAKEL DIE VOLGENDE NOODNUMMER 082 446 8946 / 0861 555 777

Signal[®] is a registered trademark of MBFi (PTY) LTD /
Signal[®] is 'n geregistreerde handelsmerk van MBFi (EDMS) BPK.



ZAR 001

Performance

WARNINGS:• **GHS Hazard Statements:**

Pictograms and Signal word	Hazard Statements	
Danger  	H314	Causes severe skin burns and eye damage.
	H411	Toxic to aquatic life with long lasting effects.

• **Relevant Substance:**

Name	CAS No.	Weight (%)
Copper sulfate	7758-99-8	1.25
Manganese sulfate	10034-96-5	6.23
Monoethylethamine (MEA)	141-43-5	17.62
Zinc oxide	1314-13-2	4.35

Although this fertilizer has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label, as well as by the method, time and accuracy of the application. Furthermore, the registration holder does not accept responsibility for damage to crops, vegetation, the environment, harm to man or animal, or lack of performance of the fertilizer concerned, due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS:• **GHS Precautionary Statements:**

P260	Do not breathe mist and spray.
P264	Wash hands thoroughly after handling.
P273	Avoid release to the environment.
P280	Wear protective gloves, protective clothing, eye protection and face protection.
P301 P330 P331	IF SWALLOWED: Rinse mouth. Do NOT induce vomiting.
P305 P351 P338	IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing.
P363	Wash contaminated clothing before reuse.
P391	Collect spillage.

- Do not eat, drink or smoke whilst mixing or applying.
- Wash hands and face before eating, drinking, smoking or using the toilet.
- Avoid spray drift onto adjacent crops, grazing, rivers and dams.
- Clean applicator and mixing containers thoroughly after use and dispose of wash water where it will not contaminate crops, grazing, rivers, dams and boreholes.
- Triple rinse the empty container: Invert the empty container over the spray or mixing tank and drain for at least 30 seconds after the flow has slowed down to dripping. Thereafter, rinse the empty container three times in succession with one quarter of the container volume fresh water and decant the rinsate into the spray or mixing tank. Puncture the triple rinsed container and dispose of via an approved collector or recycler (www.croplife.co.za). Do not bury, burn or donate the container to any other parties that may use it as a container for food or beverages.
- Do not re-use the container for any other purpose.
- Prevent contamination of food, feedstuffs, drinking water and eating utensils.

DIRECTIONS FOR USE:

- SHAKE BEFORE USING.
- Use only as indicated on this label.
- Use exclusively as fertilizer.

Row Crops:	
Foliar Applications:	Dilute 1 L/100 L water and apply to the crop (Max concentration of 3% or 3 L/100 L of water).
Soil Applications:	3-5 L/ha application.
Infurrow Applications:	1-3 L/ha application.
Tree Crops and Vines:	
Foliar Applications:	Dilute 500 ml/100 L water and apply as a full cover application (Max concentration of 2% or 2 L/100 L of water).
Soil Applications:	3-10 L/ha per application, dilute 1:1000 and apply 3-6 times through the growth periods.

Mixing: Signal® Pro should preferably be applied in the early morning or late afternoon. The rate of application is dependent on the soil fertility and reaction required. The lower rates should be used for maintenance while the higher rates will correct nutritional deficiencies.

Storage: Signal® Pro should be stored out of direct sunlight and in a dry and cool place under frost-free conditions. Keep the product locked up and out of the reach of children and animals. Avoid contamination of foodstuff or water sources.

Crops	Rate/100 L Water	Remarks
Bean: Dry & Green	500 ml Max rate of 4 L/ha	Begin application prior to pod set and repeat every 7-14 days.
Bulb Vegetables: Bulb onion, Garlic, Leek, Spring onion	500 ml Max rate of 4 L/ha	Apply at bulb initiation and repeat every 7-14 days.
Cruciferae: Broccoli, Brussels sprouts, Cabbage, Chinese cabbage, Kale, Kohlrabi, Mustard	500 ml Max rate of 3 L/ha	Apply weekly, ensuring good wetting of the entire plant beginning prior to onset of head formation.

Crops	Rate/100 L Water	Remarks
Cucurbit: Cucumber, Pumpkin, Melon, Watermelon	500 ml Max rate of 4 L/ha	Apply at flowering and repeat with 14 day intervals.
Leafy Vegetables: Chard, Chicory, Lettuce, Mung bean sprouts, Radish, Spinach, Watercress	500 ml Max rate of 3 L/ha	Apply weekly, ensuring good wetting of the entire plant beginning prior to onset of head formation.
Pea	500 ml Max rate of 4 L/ha	Apply at pod initiation and repeat every 7-14 days.
Pepper	500 ml Max rate of 4 L/ha	Begin application prior to fruit set and repeat every 7-14 days.
Pome Fruit: Apple, Pear	200-300 ml Max rate of 3 L/ha	Apply 3-4 applications from 50% flowering and repeat with 14 day intervals. For Pear: Do not combine Signal® Pro with other foliar feeds or adjuvants.
Potato	500 ml Max rate of 4 L/ha	Apply at tuber initiation and repeat every 7-14 days.
Stone Fruit: Almond, Apricot, Cherry, Nectarine, Peach, Plum, Prune	200-300 ml Max rate of 3 L/ha Small Tree (1-3) Year Max 1 L/ha Medium Tree (4-6) Year Max 2 L/ha Large Tree (7 Plus) Year Max 3 L/ha	Apply 3-4 applications from 75% petal drop and repeat at 14 day intervals. If weather conditions favour bacterial infections, additional applications can be considered. If possible, rather spray low concentration more frequently.
Tobacco	500 ml Max rate of 4 L/ha	Apply 2-3 applications bi-weekly starting at 30 cm in height.
Tomato	500 ml Max rate of 4 L/ha	Begin application prior to fruit set and repeat every 7-14 days.