

Soluble liquid fertilizer

Reg. No. B 5298 Act No. 36 of 1947 / Reg. Nr. B 5298 Wet Nr. 36 van 1947

READ THE LABEL BEFORE USE**KEEP OUT OF REACH OF CHILDREN AND ANIMALS**

Group/Groep

2

Fertilizer/Kunsmis

A WATER SOLUBLE LIQUID FERTILIZER WITH AMINO ACID CHELATED MICRO ELEMENTS, FOR FOLIAR FEEDING AND FERTIGATION.**Danger****Hazard Statements:**

- Causes serious eye damage.
- May damage fertility or the unborn child.
- May cause damage to organs.
- Very toxic to aquatic life with long lasting effects.

Precautionary Statements:

- Do not breathe mist/spray.
- Wash hands thoroughly after handling.
- Do not eat, drink, or smoke when using this product.

Composition:

Nitrogen (N)	49.5 g/kg	64.9 g/L	Manganese (Mn)*	26.0 g/kg	34.1 g/L
Sulfur (S)	15.2 g/kg	19.9 g/L	Molybdenum (Mo)	3.0 g/kg	3.9 g/L
Boron (B)	5.0 g/kg	6.5 g/L	Zinc (Zn)*	40.2 g/kg	52.6 g/L

*Amino Acid Chelated

S.G (at 20°C): 1.31

UN 3082

Registration Holder / Registrasiehouer: N Laboratories (PTY) LTD • Reg No.: 2015/165672/07
 No. 144, 2nd Avenue, Modder East Orchards, Delmas, 2210 • Tel: +27 (0) 82 738 0080 • Fax: +27 (0) 86 547 6711




Marketed and Distributed by / Bemerk en Versprei deur: MBFi (PTY) LTD • Reg No.: 2013/211882/07
 P.O. Box 1137, Delmas, Mpumalanga, 2210 • Email: orders@mbfi.co.za • Web: www.mbfi.co.za

X-Press® & GET® are registered trademarks of N Laboratories (PTY) LTD / X-Press® & GET® is geregistreerde handelsmerke van N Laboratories (EDMS) BPK.

Net Volume:
20 L

Nutrition

WARNINGS:• **GHS Hazard Statements:**

Pictograms and Signal word	Hazard Statements	
Danger   	H318	Causes serious eye damage.
	H360	May damage fertility or the unborn child.
	H371	May cause damage to organs.
	H410	Very toxic to aquatic life with long lasting effects.

• **Relevant Substance:**

Name	CAS No.	Weight (%)
Manganese sulphate	10034-96-5	8.15
Mono ethanol amine	141-43-5	15.27
Zinc borate	10361-94-1	2.85
Zinc oxide	7446-20-0	5

Although this fertilizer has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label, as well as by the method, time and accuracy of the application. Furthermore, the registration holder does not accept responsibility for damage to crops, vegetation, the environment, or harm to man or animal, or lack of performance of the fertilizer concerned, due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS:• **GHS Precautionary Statements:**

P260	Do not breathe mist/spray.
P264	Wash hands thoroughly after handling.
P270	Do not eat, drink, or smoke when using this product.
P273	Avoid release to the environment.
P280	Wear protective gloves, protective clothing, eye protection and face protection.
P305 P351 P338	IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing.
P308	If exposed or concerned: Call a POISON CENTER.

- Clean applicator and mixing containers thoroughly after use and dispose of wash water where it will not contaminate crops, grazing, rivers, dams and boreholes.
- Triple rinse the empty container: Invert the empty container over the spray or mixing tank and drain for at least 30 seconds after the flow has slowed down to dripping. Thereafter, rinse the empty container three times in succession with one quarter of the container volume fresh water and decant the rinsate into the spray or mixing tank. Puncture the triple rinsed container and dispose of via an approved collector or recycler (www.croplife.co.za). Do not bury, burn or donate the container to any other parties that may use it as a container for food or beverages.
- Do not re-use container for any other purpose.
- Prevent contamination of food, feed, drinking water and eating utensils.

DIRECTIONS FOR USE:

- SHAKE BEFORE USING.
- Use only as indicated on this label.
- Use exclusively as fertilizer.

Row Crops:	
Foliar Applications	Dilute 1 L:100 L water and apply to the crop (Max concentration of 3% or 3 L/100 L of water).
Soil Applications	3-5 L/ha application.
Infurrow Applications	1-3 L/ha application.
Tree Crops and Vines:	
Foliar Applications	Dilute 0.5 L:100 L water and apply as a full cover application (Max concentration of 2% or 2 L/100 L of water).
Soil Applications	5-10 L/ha per application, dilute 1:1000 and apply 3-6 times through the growth periods.

Mixing: X-Press® Functional Extra should preferably be applied in the early morning or late afternoon. The rate of application is dependent on the soil fertility and reaction required. The lower rates should be used for maintenance while the higher rates will correct nutritional deficiencies.

Storage: X-Press® Functional Extra should be stored out of direct sunlight and in a dry and cool place under frost-free conditions. Keep the product locked up and out of the reach of children and animals. Avoid contamination of foodstuff or water sources.

Crops	Dosage/ha	Remarks
Bean: Dry & Green	1-3 L	First application at V4-V6 growth stage and repeat at R1-R2 (start of flowering).
Berry	2-3 L	Apply at 4-6 leaf stage and repeat every 21 days.
Carrot	2-3 L	Apply at 4-6 leaf stage and repeat every 21 days.
Cherry	2-4 L	Start application at 50% flowering, repeat at fruit set, 28 days later and then as required.
Citrus	4-5 L	Start application at 50% flowering, repeat at fruit set, 28 days later and then as required.
Cruciferae	2-3 L	Apply at 4-6 leaf stage and repeat every 21 days.
Cucurbit	2-3 L	Apply at 4-6 leaf stage and repeat every 21 days.
Kiwi Fruit	2-3 L	Start application at 10 cm shoot length, repeat at pre-flowering, 3 mm berry size and 28 days later.
Lettuce	2-3 L	Apply at 4-6 leaf stage and repeat every 21 days.
Maize	1-3 L	First application at V4-V6 growth stage (15 cm shoot height), repeat if required at V8-V10 growth stage.
Nuts: Macadamia, Pecan	3-5 L	Start application at 50% flowering, repeat at fruit set, 28 days later and then as required.

Crops	Dosage/ha	Remarks
Olive	2-4 L	Start application at 50% flowering, repeat at fruit set, 28 days later and then as required.
Onion	2-3 L	Apply at 4-6 leaf stage and repeat every 21 days.
Ornamental	2-4 L	Apply every 21-28 days.
Peanut	1-3 L	First application at V4-V6 growth stage, repeat at R1-R2 (start of flowering).
Pepper	2-3 L	Apply at 4-6 leaf stage and repeat every 21 days.
Pome Fruit: Apple, Pear	3-5 L	Start application at 50% flowering, repeat at fruit set, 28 days later and then as required.
Potato	2-4 L	Apply 3-4 times per season: Start from 100% emergence (3-5 weeks) and repeat every 21-28 days.
Seedling	150 ml/100 L	Drench 10 days after germination and repeat 28 days later.
Soybean	1-3 L	First application at V4-V6 growth stage and repeat at R1-R2 (start of flowering).
Stone Fruit: Nectarine, Peach	3-5 L	Start application at 50% flowering, repeat at fruit set, 28 days later and then as required.
Strawberry	2-4 L	Apply at 4-6 leaf stage and repeat every 21 days.
Sugarcane	2-3 L	Start application when shoots are 30 cm high and repeat 28 days later.
Sunflower	1-3 L	Start application when plants are 20 cm high and repeat during flowering.
Table Grape & Vines	2-3 L	Start application at 10 cm shoot length, repeat at pre-flowering, 3 mm berry size and 28 days later.
Tobacco	2-3 L	Apply at 4-6 leaf stage and repeat every 21 days.
Tomato	2-3 L	Apply at 4-6 leaf stage and repeat every 21 days.
Turf	2-4 L	Apply every 21-28 days.
Wheat	1-3 L	First application at tillering and repeat during flag leaf stage if required.